

The universality question for noncommutative quantum field theory

Karl-Georg Schlesinger
Institute for Theoretical Physics
University of Vienna
Boltzmanngasse 5
A-1090 Vienna, Austria
e-mail: kgschles@esi.ac.at

Abstract

Present day physics rests on two main pillars: General relativity and quantum field theory. We discuss the deep and at the same time problematic interplay between these two theories. Based on an argument by Doplicher, Fredenhagen and Roberts, we propose a possible universality property for noncommutative quantum field theory in the sense that any theory of quantum gravity should involve quantum field theories on noncommutative space-times as a special limit. We propose a mathematical framework to investigate such a universality property and start the discussion of its mathematical properties. The question of its connection to string theory could be a starting point for a new perspective on string theory.

1 Introduction

The present day understanding of the physical world rests on two fundamental pillars: First, the structure of space-time which is described by general relativity (and in this way is directly linked to the gravitational force). Secondly, quantum field theory as the description of matter and all the non-gravitational forces of nature. The ongoing task to find a theory of quantum

gravity can be described as the quest for a single consistent theory, reconciling these two pillars. On the one hand, these two theories are mutually inconsistent while, on the other hand, there is deep interplay between them. Let us comment on this interplay in a little bit more detail, now.

In general relativity one can prove a remarkable theorem about black hole horizons: In all possible interactions of black holes, the total area of all black hole horizons, involved, does never decrease. This law can be formulated mathematically in a way which is completely analogous to the second law of thermodynamics with the horizon area playing the role of entropy and the surface gravity of the black hole taking the role of temperature (see e.g. [Wal]). Indeed, one can give analogous theorems for all four fundamental laws of thermodynamics. We would like to stress, here, that these are really *theorems* which can be proved in solely differential geometric terms. On this level, we have a formal analogy between two sets of laws with four laws, each. It is tempting to suppose, then, that this analogy does not come by accident and to interpret the four laws of black holes as laws for the thermodynamics of black holes. But then a problem arises: We attribute in this way a temperature to a black hole. But a non-zero temperature of an object in thermodynamics implies a non-vanishing black body radiation, emitted by this object. Hence, black holes have to emit radiation! But, by definition of a horizon, black holes can not emit anything in general relativity. So, there is a contradiction, here.

At this point, one brings quantum field theory into play. In quantum field theory, there is a spontaneous production and annihilation of pairs of particles and antiparticles from the energy of vacuum fluctuations. Now, imagine such a pair to come into existence very close to the event horizon of a black hole. Further imagine that during the lifetime of the pair one of the partners crosses the event horizon. By the laws of general relativity, this partner will never be back and the pair cannot annihilate any more. In this way, the partner outside the horizon comes into real existence and is emitted as radiation (the energy bill for turning a partner in a vacuum fluctuation into real existence is paid for by energy extraction from the black hole, i.e. the radiation really appears as emitted from the black hole). This is, very roughly speaking, the mechanism of Hawking radiation.

What we see from this is that general relativity *needs* quantum field theory to resolve the contradiction in the thermodynamics interpretation of some of its fundamental differential geometric theorems. On the other hand, the two theories are dramatically incommensurable, as can be seen e.g. from

the following observation (see [Wei]):

Suppose, we insert the vacuum energy density resulting from vacuum fluctuations into the right hand side of the Einstein equation, i.e. into the energy momentum tensor. It can be easily shown that this leads to an effective (positive) cosmological constant. Since one does not expect classical space-time concepts to make sense beyond the Planck length L , one employs a cut-off at L and in this way arrives at a finite value for the cosmological constant Λ . Now, a positive cosmological constant means that light emitted at a space-time point A and absorbed at a space-time point B acquires a red-shift, just by passing the vacuum. Assuming that light at A is emitted in the outmost violet part of the spectrum, we can ask for the distance from A to B such that at B the light has reached the borderline to the infrared in the spectrum, i.e. one can calculate a length of maximal visibility for the universe from the vacuum fluctuations of quantum field theory. The result is that when standing straight you can hardly see the floor. Hence, general relativity and quantum field theory fundamentally clash, in this example, by providing a truly dramatic contradiction to everyday experiment.

The conclusion one has to draw from this seems to be the following: On the one hand, the deep interplay between the two theories seems to indicate that general relativity and quantum field theory are viable starting points for the search for a theory of quantum gravity. On the other hand, it is clear that these two theories cannot stand just side by side in unmodified form. This is precisely the perspective, we will take in this paper.

The second fundamental input to this paper comes from the results of [DFR]. We have already remarked above that one does generally not believe that a classical description of space-time does hold at or beyond the Planck length. It has been argued for decades that a foam-like structure of space-time should appear at this scale. In [DFR] this has been put into a rigorous form: It is shown there that the creation of black holes from vacuum fluctuations at the Planck scale leads to uncertainty relations for space-time coordinates. Since one knows that uncertainty relations derive from noncommuting variables in quantum mechanics, this leads to the conclusion that space-time should be described by a noncommutative space in a theory of quantum gravity. Accepting the requirement for such uncertainty relations of space-time coordinates as *universal*, i.e. not linked to a specific theory of quantum gravity but being a prerequisite of any such theory, one arrives - together with the considerations from above - at the following postulate:

Any theory of quantum gravity should at least have a limit in the form of a formulation in terms of quantum field theories on noncommutative space-times.

We have required a limit, only, since one can imagine that there could exist theories which incorporate a theory of quantum gravity but go beyond providing only this. We will call this postulate the *universality postulate*, in the sequel. The aim of the present paper is to elaborate on this postulate. First, we try to give a model for a mathematical formulation of this postulate in section 2. Secondly, in section 3 we undertake to derive some of the consequences of such a mathematical formulation. As we will discuss in the next section, this can also be seen as an Ansatz for a quite different approach to string theory.

We want to stress that this is not a research paper in a proper sense. Rather it is intended as giving one possible motivation for later technical work ([Sch 2006] and planned follow-up work) and taking a short view from a broader perspective.

Remark 1 *For a paper which also follows the idea of a universality of non-commutativity but investigates conclusions in a quite different direction, see [DMK].*

2 A mathematical framework

Consider a smooth n -dimensional manifold M endowed with a Poisson bracket. In [Kon], it was shown that the deformation theory of such a manifold is described by a universal formula. Moreover, this formula derives from a physical system given by a two dimensional field theory (this description in terms of the Poisson sigma model introduced in [I], [SchSt] was made explicit in [CF]).

In analogy, one can consider the following situation: Suppose we would be given a mathematical description of quantum field theory together with its deformations to quantum field theories on noncommutative space-times (henceforth noncommutative quantum field theories or ncQFTs, for short). Of course, even for the classical case of QFTs on usual Minkowski space a mathematical formulation is a widely open problem and is considered as one

of the deepest questions of 21st century mathematics. We could introduce, then, a hypothetical category $\mathcal{A}$ (possibly with some extra structure, see below) of all (commutative and noncommutative) QFTs. We can ask, then, if $\mathcal{A}$ is related to some physical system, inheriting all the symmetry and deformation properties of objects in $\mathcal{A}$ (much the same way as [CF] gives the physical system for deformation theory of Poisson manifolds). Even more concrete, we can ask if such a system - if it exists - is related to string- or M -theory, in this way, in principle, opening up the possibility to approach string theory from a completely different viewpoint by posing the question if it might be related to a universal deformation theory for quantum field theories (and, hence, be universal in the sense of the above universality postulate).

Given such a category $\mathcal{A}$, the universality postulate of the introduction would translate to the postulate that any consistent theory of quantum gravity should have a formulation in terms of a family of categories $\mathcal{B}(t)$ such that a limit

$$\mathcal{B}(t) \rightarrow_{t \rightarrow 0} \mathcal{A}$$

exists. Here, t is a deformation parameter ($t \in \mathbb{R}$) and the family $\mathcal{B}(t)$ and the limit are understood in the sense of formal deformations.

As we have mentioned, even the formulation of the category $\mathcal{A}$ might be out of reach for a long time to come. We therefore proceed as follows: We suggest a simpler model for an analogous situation and study some of its properties. Even in this case, the category $\mathcal{A}$ will remain a partly conjectural object and one should read the present paper more as a suggestion for future work rather than a statement of final results.

Let us pass to the special case of two dimensional conformal field theories. A central structural element of these theories is the so called operator product expansion (OPE). The OPE has a rigorous mathematical formulation in terms of vertex algebras (also called chiral algebras in the more abstract, coordinate independent, formulation of [BD]). For the case of two dimensional rational conformal field theories (RCFTs), one can show that the structure of a vertex algebra completely determines these theories (see [BPZ], [Wit], [FFFS]), i.e. for the highly restricted class of RCFTs a completely rigorous mathematical formulation is available. One possible formulation is given in terms of certain inner Frobenius algebras of braided monoidal categories (see [FS], [FRS 2001], [FRS 2003], [SFR]).

The deformation theory of vertex algebras has been studied in special cases and shown e.g. to correspond in some examples to massive deforma-

tions of RCFTs (see e.g. [BK], [FL], [Fre], [FR], [Kli], [Tam]). In an abstract setting (extracting only the general properties of the product structures in a vertex algebra), we have studied a general deformation theory of vertex algebras in [GS]. We have shown there that the deformation theory is described by a system of differential equations (replacing the Maurer-Cartan equation of the deformation theory of usual associative algebras) and that this system can formally be derived from an action principle.

In the next few pages, we will collect some of the properties of the deformation complex of [GS]. We will suggest, then, a model for the category $\mathcal{A}$ in terms of these deformation complexes.

We now want to discuss some of the structural properties of the deformation complex of a quantum vertex algebra as described by the action (18) of [GS]. We first restrict to the case where, only the structure of one monoidal category is deformed, i.e. to

$$\begin{aligned} S \sim & \int \{ f \circ d_{\bullet} f + \frac{2}{3} f \circ f \circ f + g \circ d_{\otimes} g + \frac{2}{3} g \circ g \circ g \\ & + \lambda \circ^2 [d_{\otimes}^{\otimes 2} f - d_{\bullet}^{\otimes 2} g + f \otimes f - g \bullet g - (f \circ^2 g) + (g \circ^2 f) \\ & - Comp(f, g, g) + Comp(g, f, f)] \} \end{aligned}$$

Remember that on the usual Hochschild complex of associative algebras one has the structure of a graded homotopy Lie algebra given by the differential and the Gerstenhaber bracket. Together with the wedge product this combines into a homotopy Gerstenhaber algebra. In the case of a monoidal category, we have a more complicated structure on the deformation complex, consisting of the following elements:

- As in the case of the Hochschild complex, we have the composition $\circ$, the graded commutator of $\circ$ giving the Gerstenhaber bracket.
- We have two different differentials $d_{\bullet}$ and $d_{\otimes}$ plus their liftings $d_{\bullet}^{\otimes 2}$ and $d_{\otimes}^{\otimes 2}$ to the twofold tensor product.
- We have the graded “curvature” tensor $Comp$.
- The products $\bullet$ and $\otimes$ themselves are represented on the deformation complex.

- Observe that, in contrast to the Hochschild complex, all these structures are not defined with respect to a single vector space but with respect to the whole collection of Hom -sets.

Remark 2 *The two differentials $d_\bullet$ and $d_\otimes$ are not compatible in the sense of a double complex. Actually, the compatibility condition does not make sense on the deformation complex of a monoidal category because by the composability requirements we have imposed on the deformations of the composition in [GS], the two expressions $d_\bullet d_\otimes f$ and $d_\otimes d_\bullet f$ are not well defined at the same time.*

The fact that $\bullet$ and $\otimes$ themselves appear on the deformation complex, again, is a very unusual property. It means that the full structure of the monoidal category can be reconstructed from the deformation complex. The monoidal category does not only determine its deformation complex but also vice versa. This also remains true in the full setting of the action (18) of [GS]: Besides the structure of both monoidal categories, also the functor $\mathcal{F}$ appears on the deformation complex and can therefore be reconstructed from the complex. The following question therefore arises: Can the additional structure on the complex - appearing beyond the monoidal category structures of $\mathcal{C}$ and $\mathcal{M}$ and the monoidal functor $\mathcal{F}$ - be interpreted as additional structure on the triple $(\mathcal{M}, \mathcal{C}, \mathcal{F})$? In other words: Can one start with categories $\mathcal{C}$ and $\mathcal{M}$ which carry more structure than the one of a monoidal category and a functor $\mathcal{F}$ adapted to this additional structure such that one gets a full duality between the triple $(\mathcal{M}, \mathcal{C}, \mathcal{F})$ and its deformation complex? If yes, is this additional structure related to the additional structure which should be carried by a quantum vertex algebra?

In a first approach toward these questions, we restrict, again, to the setting of only one monoidal category $\mathcal{C}$ as described by the action given above. We recall that in [BD] a pseudotensor category is defined by the following data (where $\mathcal{S}$ denotes the category of finite sets and surjective maps): A class $\mathcal{M}$ of objects together with

1.) For any $I \in \mathcal{S}$, an I -family of objects $L_i \in \mathcal{M}$, $i \in I$, and an object $M \in \mathcal{M}$, there exists the set $P_I^{\mathcal{M}}(\{L_i\}, M)$ of I -operations.

2.) For any surjective map $\pi : J \twoheadrightarrow I$ in $\mathcal{S}$, families of objects $\{L_i\}_{i \in I}$, $\{K_j\}_{j \in J}$ and an object M , there exists a composition map

$$P_I^{\mathcal{M}}(\{L_i\}, M) \times \prod_{i \in I} P_{J_i}^{\mathcal{M}}(\{K_j\}, L_i) \rightarrow P_J^{\mathcal{M}}(\{K_j\}, M)$$

with

$$(\varphi, (\psi_i)) \longmapsto \varphi(\psi_i)$$

and $j \in J_i$ in $P_{J_i}^{\mathcal{M}}(\{K_j\}, L_i)$ where $J_i = \pi^{-1}(i)$.

The composition is assumed to be associative and the existence of units is assumed (see [BD]). Recall that a pseudotensor category with one object is an operad. A representable pseudotensor category is equivalent to a usual category $\mathcal{M}$ together with functors

$$\otimes_I : \mathcal{M}^I \rightarrow \mathcal{M}$$

for any $I \in \mathcal{S}$ and a natural compatibility morphism

$$\epsilon_\pi : \otimes_J K_j \rightarrow \otimes_I (\otimes_{J_i} K_j)$$

- for any surjective map $\pi : J \twoheadrightarrow I$ - satisfying certain naturality conditions (see [BD]). If all ϵ_π are isomorphisms, the pseudotensor category reduces to a symmetric monoidal category.

Now, observe that the deformation complex of a monoidal category $\mathcal{C}$ is given by the multilinear maps

$$Morph(\mathcal{C}^k) \rightarrow Morph(\mathcal{C}) \tag{1}$$

for any $k \in \mathbb{N}$ (where the multilinear maps are understood as those which are multilinear locally on the *Hom*-sets). Considering the set of such multilinear maps for fixed k as the set of k -operations, one can define a composition on the deformation complex in complete analogy to the composition in a pseudotensor category. One has a kind of pseudotensor 2-category with one object in this way (i.e. a kind of categorical operad). We write “a kind of” because on the one hand, all the maps in (1) live on the category $\mathcal{C}$ but on the other these maps are *not* functors. If we want to stress this fact, we will speak of a *pseudotensor semi-2-category*.

It is a straightforward consequence that the tensor $Comp$ is induced from this pseudotensor composition as a special case. So, we get a kind of categorical version of the pseudotensor structure of [BD] as part of the additional structure on the monoidal category $\mathcal{C}$, given by the deformation complex.

We phrase the following question:

Question: Can the full additional structure, given by the deformation complex, be understood as a (kind of categorical) extension of the chiral algebra structure of [BD] (possibly after extending the consideration to the full triple $(\mathcal{M}, \mathcal{C}, \mathcal{F})$)? Does a duality between the triple $(\mathcal{M}, \mathcal{C}, \mathcal{F})$ and its deformation complex hold?

Suppose the answer to the above question would be in the affirmative. The general noncommutative deformation of a RCFT could then be understood as being given by what a category theorist would probably call a 2-chiral algebra. In the sequel, let us assume this to be true. Whatever the precise definition of a 2-chiral algebra will be (we assume it to resemble the structure of the deformation complex, described above), a 2-chiral algebra will certainly be given by a bicategory with some extra structure. It is this lack of a precise definition of a 2-chiral algebra and our ignorance of the answer to the above question which makes even the simple toy model we suggest for $\mathcal{A}$ a largely speculative object. Assuming that we have bicategories with some extra structure, all 2-chiral algebras should ensemble into a tricategory (see [GPS] for the definition of tricategories or the introduction included in [Lau]). This tricategory will be our toy model for $\mathcal{A}$.

3 Some properties of $\mathcal{A}$

Let us discuss, now, some of the possible properties of $\mathcal{A}$. This discussion will, again, be largely speculative. Remember that we understand this paper, only, as a motivation for later technical work.

In formal deformation theory, one can often distinguish between the deformations of an algebraic structure inside the given category and the - more general - quantum deformations which usually lead to a categorification of the structure. E.g. one has deformations of Lie algebras - or their universal envelopes - as such and quantum deformations inside the more general category of Hopf algebras. In [Sch 2001], we have given an argument that

certain monoidal bicategories should not have nontrivial quantum deformations, i.e. all deformations should remain in the given category of these special monoidal bicategories. We called this ultrarigidity.

Let us speculatively assume that ultrarigidity also holds true for 2-chiral algebras (observe that the structure on the deformation complex, described in the previous section, seems to be a generalization of a class of monoidal bicategories). It follows, then, that all deformations of 2-chiral algebras remain inside $\mathcal{A}$. If there would be a deformation of a 2-chiral algebra to an algebraic structure outside $\mathcal{A}$, this should correspond to a deformation of $\mathcal{A}$ itself. Conversely, a deformation of $\mathcal{A}$ should correspond to a deformation of the notion of 2-chiral algebra, i.e to a deformation of the objects of $\mathcal{A}$. If ultrarigidity holds for 2-chiral algebras, this implies that any deformation of $\mathcal{A}$ should correspond to a functor $\mathcal{F}$ of tricategories,

$$\mathcal{F} : \mathcal{A} \rightarrow \mathcal{A} \tag{2}$$

(If it would not be a true functor of tricategories, this would mean that the morphism structure of $\mathcal{A}$ would be changed in the process of deformation. But this, in turn, would mean that the notion of 2-chiral algebra has been deformed, i.e we would have deformations of 2-chiral algebras which do not stay inside $\mathcal{A}$, contradicting ultrarigidity. Hence, $\mathcal{F}$ has to be a functor of tricategories, once we have assumed ultrarigidity.). Conversely, if any deformation of $\mathcal{A}$ would lead to a functor (1), this should imply ultrarigidity. Observe that $\mathcal{F}$ does not have to be invertible or even an equivalence of tricategories. So, there are still two possibilities for deformations: On the one hand, the trivial deformations corresponding to equivalences of $\mathcal{A}$ and, on the other hand, the nontrivial deformations which can e.g. lead to proper sub-tricategories of $\mathcal{A}$. So, there is still room for quantization of 2-chiral algebras in the sense of nontrivial deformations of $\mathcal{A}$ but these would be 2-chiral algebras, again. Especially, ultrarigidity would imply that quantization of 2-chiral algebras has to be functorial.

So, assuming that - in the narrow sense of our toy model - commutative and noncommutative quantum field theories would be described by 2-chiral algebras, all the important questions how such quantum field theories ensemble into families, i.e. how a single quantum field theory is deformed in a family, how they are renormalized (as a special case of deformation theory in a family of quantum field theories), etc. should have a counterpart in the deformation theory of $\mathcal{A}$. Especially, with a detailed knowledge about

the correct definition of 2-chiral algebras and the tricategory $\mathcal{A}$, it should be possible to prove ultrarigidity for 2-chiral algebras from the deformation theory of $\mathcal{A}$ by checking for the existence of the functors (1). As a very first step, we will study the deformation theory of tricategories in the forthcoming paper [Sch 2006].

Remark 3 *Assuming the universality postulate from above, proving ultrarigidity from the deformation theory of $\mathcal{A}$ would imply that $\mathcal{A}$ is singled out as providing the only consistent framework for a quantum theory of gravity. A possible physical system related to $\mathcal{A}$ (see below) would in this way be singled out on the basis of the universality postulate and mathematical - deformation theoretic - arguments.*

There are two important questions, then:

- Does the deformation theory of $\mathcal{A}$ fully determine the structure of $\mathcal{A}$, much the same way we have discussed this for the deformation theory of vertex algebras, in the previous section?
- Does the deformation theory of $\mathcal{A}$ (deformation equations, replacing the Maurer-Cartan equation, cohomology, etc.) arise from a physical system?

Since chiral algebras correspond to two dimensional quantum field theories, one would suspect - in the spirit of the dimensional ladder and categorification - that 2-chiral algebras should correspond to field theories on the three dimensional world volume of a membrane (possibly topological, since usual chiral algebras correspond to RCFTs). If this would be true, it would naturally fit in with the idea of [GMSS] that noncommutative quantum field theories should, in a very general sense, arise from open membrane systems. So, it seems natural to sharpen the second of the above questions to the question if the deformation theory of $\mathcal{A}$ could arise from (topological) M -theory. If the answer would be in the affirmative, this would open up the possibility to study M -theory as the universal theory related to the deformation theory of (noncommutative) quantum field theories (much the same way as [CF])

gives the universal theory related to the deformation theory of Poisson manifolds). It will need rigorous technical work to judge if such a suggestion is anything more than pure speculation.

Remark 4 *For a different approach which also suggests that string theory might have universality properties as a theory ruling the behaviour of QFTs, see [Vaf].*

Acknowledgments: I would like to thank H. Grosse and V. Mathai for discussions on or related to the material of this paper.

References

- [BD] A. Beilinson, V. Drinfeld, *Chiral algebras*, American Mathematical Society, Providence 2004.
- [BK] B. Bakalov, V. Kac, *Field algebras*, math.QA/0204282.
- [BPZ] A. A. Belavin, A. M. Polyakov, A. B. Zamolodchikov, *Infinite conformal symmetries in two dimensional quantum field theory*, Nucl. Phys. B241 (1984), 333-380.
- [CF] A. Cattaneo, G. Felder, *A path integral approach to the Kontsevich quantization formula*, math/9902090.
- [DFR] S. Doplicher, K. Fredenhagen, J. E. Roberts, *The quantum structure of spacetime at the Planck scale and quantum fields*, hep-th/0303037.
- [DMK] M. Dubois-Violette, J. Madore, R. Kerner, *Shadow of noncommutativity*, q-alg/9702030v2.
- [FFFS] G. Felder, J. Fröhlich, J. Fuchs, C. Schweigert, *Conformal boundary conditions and three-dimensional topological field theory*, hep-th/9909140.

- [FL] G. Felder, A. LeClair, *Restricted quantum affine symmetry at perturbed minimal conformal models*, hep-th/9109019.
- [Fre] E. Frenkel, *Affine Kac-Moody algebras, integrable systems and their deformations*, math.QA/0305216.
- [FR] E. Frenkel, N. Reshetikhin, *Towards deformed chiral algebras*, q-alg/9706023v2.
- [FRS 2001] J. Fuchs, I. Runkel, C. Schweigert, *Conformal correlation functions, Frobenius algebras and triangulations*, hep-th/0110133v2.
- [FRS 2003] J. Fuchs, I. Runkel, C. Schweigert, *Boundaries, defects and Frobenius algebras*, hep-th/0302200.
- [FS] J. Fuchs, C. Schweigert, *Category theory for conformal boundary conditions*, math.CT/0106050v2.
- [GMSS] R. Gopakumar, S. Minwalla, N. Seiberg, A. Strominger, *OM theory in diverse dimensions*, hep-th/0006062.
- [I] N. Ikeda, *Two-dimensional gravity and nonlinear gauge theory*, Ann. Phys. 235 (1994), 435-464.
- [Kli] C. Klimcik, *Quasitriangular WZW model*, hep-th/0103118.
- [Kon] M. Kontsevich, *Deformation quantization of Poisson manifolds I*, math/9709180.
- [Lau] A. Lauda, *Frobenius algebras and planar open string topological field theories*, math.QA/0508349
- [Mac] S. Mac Lane, *Categories for the working mathematician*, Springer, Berlin 1971.
- [GPS] R. Gordon, A. J. Power, R. Street, *Coherence for tricategories*, AMS, Providence 1995.
- [GS] H. Grosse, K.-G. Schlesinger, *On deformation theory of quantum vertex algebras*, hep-th/0508225.

- [Sch 2001] K.-G. Schlesinger, *A quantum analogue of the Grothendieck-Teichmüller group*, J. Phys. A: Math. Gen. 35 (2002), 10189-10196, math.QA/0104275.
- [Sch 2006] K.-G. Schlesinger, *Does the fivebrane have a nonclassical BV-structure?*, in preparation.
- [SchSt] P. Schaller, T. Strobl, *Poisson structure induced (topological) field theories*, Modern Phys. Lett. A9 (1994), no. 33, 3129-3136.
- [SFR] C. Schweigert, J. Fuchs, I. Runkel, *Categorification and correlation functions in conformal field theory*, math.CT/0602079.
- [Tam] D. Tamarkin, *Deformations of chiral algebras*, math.QA/0304211.
- [Vaf] C. Vafa, *The string landscape and the swampland*, hep-th/0509212v2.
- [Wal] R. M. Wald, *General relativity*, University of Chicago Press, Chicago 1984.
- [Wei] S. Weinberg, *The cosmological constant problem*, Rev. Mod. Phys. 61 (1989), 1-23.
- [Wit] E. Witten, *Quantum field theory and the Jones polynomial*, Comm. Math. Phys. 121 (1989), no. 3, 351-399.